

Tuberculosis

TB – the disease, its treatment and prevention



TUBERCULOSIS



TB (tuberculosis) is an infectious disease that usually affects the lungs, although it can affect any part of the body.

About 150 years ago, it caused about one in eight of all deaths in the UK, but by the 1980s, with better housing and nutrition and effective treatments, it had become uncommon in the UK with 5745 cases in 1987.

However, TB had not been wiped out completely. Over the last 20 years numbers in the UK have been rising slowly. About 7000 people now get TB each year – just over one person in every 10,000 of the population.

TB is not easily caught – you have to be in close and lengthy contact with someone with TB (for example, living in the same household) – but everybody should be aware of the symptoms of the disease so they can seek treatment as soon as possible.

TB is curable with a course of special antibiotics.

How is TB spread and am I likely to get infected?

TB can only be caught directly from someone with infectious TB in their lungs or throat. Although TB is spread through the air when people who have the disease cough or sneeze, it takes close and lengthy contact with an infectious person to catch the disease. So it's unlikely that you'll catch TB on the bus or tube, for example.

Not everyone with TB of the lungs is infectious, and as long as they are taking the proper treatment most people that were infectious become non-infectious pretty quickly – generally after about two weeks – **as long as they are taking the proper treatment.**

While anyone can catch TB, some groups of people are more at risk than others. These include people who:

- have lived in the same household – or been in close and lengthy contact – with someone with infectious TB
- are living in unhealthy or over-crowded conditions, including those who are homeless or sleeping rough
- have lived, worked or stayed for a long time in a country with a high rate of TB, such as south-east Asia, sub-Saharan Africa and some countries in eastern Europe
- may have been exposed to TB in their youth when the disease was more common in this country
- are the children of parents whose country of origin has a high rate of TB
- have been in prison
- are unable to fight off infection (immunosuppressed) due to illness (e.g. HIV infection) or treatment
- are addicted to drugs or misuse alcohol
- do not eat enough to stay healthy.

How will I know if I've got TB?

The most common symptoms of TB include:

- a persistent cough that gets progressively worse over several weeks
- loss of weight for no obvious reason
- fever and heavy night sweats
- a general and unusual sense of tiredness and being unwell
- coughing up blood.

All these may also be signs of other problems but if you have them and are worried, talk to a doctor or nurse at your local surgery or clinic, or go to an NHS walk-in centre or phone NHS Direct on 0845 4647.

(For patients' safety, all calls to NHS Direct are recorded. Calls are charged at the same rate as local calls.)

If you are a close contact of someone who has been diagnosed with TB and

there is a risk you may have the infection, you will be offered a check-up at a special TB clinic.

If I have TB, can I be cured?

Yes, TB can be treated with special antibiotics. Once treatment starts, you will begin to feel better after about two to four weeks. But the treatment has to continue for at least six months. It is vitally important to complete the whole course of antibiotics to cure the TB. If you don't, the TB may return in a form that is resistant to the usual drugs and much more difficult to treat. And you can die from this more serious form of the infection. Tell your family and

If TB is not treated properly, it may lead to death.



I thought TB was prevented by a vaccine?

There is a vaccine (BCG) that has been in use for many years to help protect against TB. BCG works best to prevent the more serious forms of TB in children. However, the BCG vaccine does not prevent TB in all cases so you still need to know the signs and symptoms of TB.

Who is offered BCG vaccination?

BCG vaccine is offered to:

- infants born or living in areas where the incidence of TB is 40/100,000 of the population or greater
- infants whose parents or grandparents were born in or come from a country with a TB incidence of 40/100,000 of the population or greater
- previously unvaccinated new immigrants from high prevalence TB countries.

How else can TB be prevented?

The most important and effective way to prevent TB spreading in this country is to diagnose people with the disease as soon as possible and make sure they have a full course of correct treatment. That is why it is so important to know about the disease.

I'm going abroad, do I need a BCG injection?

If you are going to be visiting, living or working in a country with a high rate of TB for more than one month, it's important that you are protected from TB. Ask for advice at your doctor's surgery or clinic.

TB IN NUMBERS FACTFILE

- TB has been around for at least 4000 years – Egyptian mummies show signs of the disease.
- Around 2 billion people (about one third of the world's population) are infected with TB – and are therefore at risk of developing active TB (TB disease) at some time in their life.
- The World Health Organization (WHO) estimates that across the world, almost 9 million people develop TB disease every year.
- TB also causes around 2 million deaths every year. Alongside malaria and HIV it is one of the greatest infectious disease challenges in the world today.
- In sub-Saharan Africa, the HIV epidemic is making people much more susceptible to developing TB disease.
- In the UK as a whole, TB has increased by 27 per cent since the 1980s, from 5745 to 7300 cases a year.
- In London, the number of TB cases has doubled since the 1980s, accounting for almost 3000 cases each year.
- In the UK, because of good access to effective services, death from TB is rare.

TB – common concerns

Most people living in the UK will never encounter a case of TB. While it is important that people are aware of TB symptoms, and know how TB is spread and treated, it is also important that people understand the real risk to themselves and their families.

Public transport and enclosed public places

Using public transport and going about your normal daily business does not put you at increased risk of getting TB.

TB in schools

This is very rare but can cause great anxiety. Children with TB hardly ever spread the disease. Children catch TB from adults with TB of the lungs. If a child in school were found to have TB then screening will be organised to make sure no one else is infected and to identify the source of infection. The source is usually a close family member or someone from the same household. If an adult member of staff were found to have infectious TB

then children would be screened following local public health advice.

Why has the schools BCG programme stopped if TB is increasing?

The pattern of TB has changed considerably since the schools BCG programme was introduced in 1953 and there is no longer a justification to vaccinate children at very low risk from TB. The new policy offers BCG to those people who are most likely to benefit from vaccination as early in life as possible when the vaccine is most effective. (See the Department of Health TB factsheet for more information).

TB and immigration

It is true that over half of the people diagnosed with TB in this country were born abroad but immigration alone does not explain the recent rise in TB cases. Around forty per cent of people born overseas who develop TB in this country have lived here for over ten years. The most important action to safeguard public health is to ensure that all TB cases are diagnosed promptly and fully treated.

More information

For more information on protecting yourself, your family and friends against TB, you can contact:

NHS Direct on 0845 4647* or talk to your doctor or TB Alert

* For patients' safety, all calls to NHS Direct are recorded. Calls are charged at the same rate as local calls.

TB Alert is a charity dedicated to raising awareness about TB and fighting TB worldwide.

TB Alert

22 Tiverton Road

London

NW10 3HL

Phone: 0845 456 0995

e-mail: info@tbalert.org

Reg Charity No: 1071886

You can also visit

www.immunisation.nhs.uk

www.dh.gov.uk

www.tbalert.org

www.hpa.org.uk

NHS Direct Interactive – a health information service available on digital satellite TV by pressing the interactive button on the remote control.

Kjo fletëpalosje është e disponueshme në gjuhën shqipe në adresat e mëposhtme në internet. (Albanian)

এই পত্রিকাটি নিচের ওয়েব ঠিকানায় বাংলায় পাওয়া যায় (Bengali)

這本中文繁體字版小冊子載於下列網址。 (Chinese)

این بروشور به زبان فارسی در آدرس وب زیر موجود است. (Farsi)

Cette brochure est disponible en français sur les sites Web ci-dessous (French)

Αυτό το φυλλάδιο διατίθεται στα Ελληνικά στις παρακάτω διευθύνσεις στο Διαδίκτυο (Greek)

આ લીફલેટ ગુજરાતીમાં નીચે જણાવેલા સરનામાંઓ પર ઉપલબ્ધ છે. (Gujarati)

Questo depliant è disponibile in italiano sui siti web sotto indicati (Italian)

دغه رساله په پښتو ژبه په لاندې ویب شتهد. (Pashto)

Niniejsza ulotka jest dostępna w języku polskim pod adresami internetowymi wymienionymi poniżej (Polish)

O folheto está disponível em português no endereço da Internet abaixo (Portuguese)

ਇਹ ਲੀਫਲੈਟ ਹੇਠਾਂ ਦਿੱਤੇ ਗਏ ਵੈੱਬ ਪੱਤਿਆਂ 'ਤੇ ਪੰਜਾਬੀ ਵਿਚ ਉਪਲਬਧ ਹੈ। (Punjabi)

Acest pliant poate fi găsit în limba română la adresa web de mai jos (Romanian)

Xaashidaan yar waxaad ku heli kartaa af Soomaali cinwaanka internet'ka ee hoos ku qoran (Somali)

وه رگپراوي سۆرانیی ئه م ناملیکه یه له ریگای ئه م ماڵپه ره ی خواره وه ده ست ده که وئ. (Sorani)

Este folleto está disponible en español en las direcciones de sitio web de más abajo (Spanish)

கீழ்க்கண்ட வலைதள முகவரிகளில் இந்தத் துண்டுப் பிரசுரம் தமிழில் கிடைக்கிறது. (Tamil)

Bu kitapçık Türkçe olarak aşağıdaki web adresinde mevcuttur (Turkish)

یہ پرچہ اردو میں درج ذیل ویب پتوں پر دستیاب ہے۔ (Urdu)

Văn bản in rời này có bản bằng tiếng Việt tại địa chỉ web dưới đây. (Vietnamese)

This leaflet is available in the above languages on the website www.immunisation.nhs.uk

Tuberculosis

TB – the disease, its treatment
and prevention

Further copies of this leaflet
can be obtained from:
Department of Health Publications
Phone: 08701 555 455
e-mail: dh@prolog.uk.com
Textphone (for minicom users)
0870 0102 870 for hard of hearing
8am to 6pm Mon-Fri

For more information on
immunisation visit our website at
www.immunisation.nhs.uk



Immunisation Information